Appendix A: Health and Wellbeing Performance Metrics 2015/16

| | Updated indictors are bolded | Frequency | Latest Period of Availability | Previous Available Period (Lewisham | Latest Available Period (Lewisham) | Lon | Eng | England Benchmark | | Data Source |
|----------|---|----------------------------|-------------------------------------|--|---|---------------|--------------|----------------------|----------|--|
| | Overarching Indicators | | | · · | | | | | | |
| 1a | Life Expectancy at Birth (Male)(yrs) | Annual | 2011-13 | 78.2 | 78.7 | 80.0 | 79.4 | sig low | | ONS |
| | Life Expectancy at Birth (Female)(yrs) | Annual | 2011-13 | 82.6 | 83.0 | 84.1 | 83.1 | similar | | ONS |
| | Children in poverty (%) Under 75 from CVD mortality (DSR) | Annual Annual | 2012 2011-13 | 30.5 91.0 | 27.7 87.4 | 23.7 80.1 | 19.2 78.2 | sig high sig high | | CHIMAT Profile 2015 NHSIC - P00400/ PHOF 4.04i |
| | Potential Years of Life Lost (PYLL) from causes considered amenable to healthcare (DSR) | Annual | 2014 | - | 2212.6 | - | 2064.5 | sig high | <u> </u> | NHSOF 1A - ONS (CCG 1.1 DSR)- P01559 |
| | Slope index of inequality in life expectancy at birth within English local authorities, based on local deprivation deciles within each area (Male) | Annual | 2011-13 | 6.6 | 7.7 | - | - | - | 1 | PHOF 0.2iii |
| 5h | Slope index of inequality in life expectancy at birth within English local authorities, based on local deprivation deciles within each area (Female) | Annual | 2011-13 | 6.6 | 6.6 | - | - | - | | PHOF0.2iii |
| | Infant Mortality (%) | Annual | 2011-13 | 4.9 | 4.6 | 3.8 | 4.0 | similar | | P00723/CHIMAT Profile 2014 |
| | Low Birth Weight of all babies (%) | Annual | 2013 | 8.4 | 7.8 | 7.9 | 7.4 | sig high | - I | P00455/CHIMAT Profle 2014 |
| | Proportion of people using social care who receive self-directed support, and those receiving direct | Annual | 2013-14 | 55.5 | 69.4 | 67.5 | 62.1 | _ | | ASCOF(1C)- NHSIC -P01509 |
| | payments Average number of delayed transfers of care on a particular day taken over the year (Crude rate per | Annual | 2013-14 | 4.9 | 4.6 | | | | | ASCOF 2C- NHSIC - P01516 |
| | 100,000 adult population) | | | | | 6.8 | 9.6 | - | _ | |
| | Total Days of Delay (crude rate per 100,000) Priority Objective 1: Achieving a Healthy Weight | Annual | 2014-15 | 1418.7 | 1570.0 | - | - | - | <u> </u> | BCF - Local Data - LPI264 (PPLUS) |
| | Excess weight in Adults (%) | Annual | 2012-13 | _ | 61.2 | 57.3 | 63.8 | similar | | PHOF 2.12 |
| | Excess weight in Children - Reception Year (%) | Annual | 2013-14 | 25.0 | 24.6 | 23.1 | 22.5 | sig high | | PHOF 2.06 |
| 12b | Excess Weight in Children- Year 6 (%) | Annual | 2013-14 | 38.3 | 39.3 | 37.6 | 33.5 | sig high | | PH NCMP Profiles |
| | Breastfeeding Prevalence 6-8 weeks(%) | Qtr | 2015-16 Q1 2014 | 73.4 | 73.9 | | 45.2 | similar | | NHS ENGLAND PHOF 2.13i (Active People Survey) |
| | % of physically active and inactive adults - Active adults % of physically active and inactive adults - Inactive adults | Annual Annual | 2014 | 58.1 25.1 | 57.1 27.5 | 56.2 27.0 | 56.0 27.7 | similar similar | | PHOF 2.13i (Active People Survey) PHOF 2.13ii (Active People Survey) |
| 140 | Priority Objective 2: Increasing the number of people who survive colorectal, brea | | | | 27.3 | 27.0 | 27.7 | Silling | - | THOI 2.15H (Netive I copie survey) |
| 15a | Cancer screening coverage - breast cancer (%) | Annual | 2014 | 66.0 | 65.0 | 68.9 | 75.9 | sig low | - | PHOF 2.20i |
| 15b | Cancer screening coverage - cervical cancer(%) | Annual | 2014 | 72.4 | 73.7 | 70.3 | 74.2 | sig low | | PHOF 2.20ii |
| 15c | Cancer screening coverage - bowel cancer (%) | Qtr - Local | 2014/15 Q1 | - | 46.4 | - | - | - | - | London Bowel Screening Hub (Local data available for 6 S.E Lon Boroughs) - 49.4% (S.E.L avg) |
| 16 | Early diagnosis of cancer (%) | Annual | 2013 | 39.9 | 45.6 | 43.1 | 45.7 | - | | PHOF 2.19 – experimental statistics |
| | Two week wait referrals (number per 100,000 population) | Annual | 2013-14 | 2273 | 2614 | - | 2399 | - | | Cancer Toolkit GP Profiles |
| | Under 75 mortality from all cancers (DSR) | Annual | 2011-13 | 159.9 | 159.2 | 136.5 | 144.4 | sig high | Ų. | NHSIC - P00381/ PHOF 4.05i |
| | Priority Objective 3: Improving Immunisation Uptake | I _ | T | | | T | | | | |
| | Uptake of the first dose of Measles Mumps and Rubella vaccine (MMR1) at two years of age | Qtr | 2015-16 Q1 | 90.0 | 87.5 | 86.7 | 92.1 | low | | PHOF 3.03viii/ Local Imms Cover Data |
| \vdash | Uptake of the second dose of Measles Mumps and Rubella Vaccine (MMR2) at five years of age | Qtr | 2015-16 Q1 | 71.0 | 70.0 | 80.4 | 88.5 | low | | Local Immunisation cover data |
| | Uptake of the third dose of Diptheria vaccine (D3) at one year of age Uptake of the fourth dose of Diphtheria vaccine (D4) at five years of age | Qtr Qtr | 2015-16 Q1 2015-16 Q1 | 92.2 83.5 | 91.0 81.5 | 90.0 78.5 | 93.6 87.7 | low | | Local Immunisation cover data Local Immunisation cover data |
| - | Uptake of Human Papilloma Virus (HPV) vaccine in girls in Year 8 in Lewisham Schools | Annual | 2013-14 | 84.8 | 82.9 | 80.0 | 86.7 | sig low | 1 | PHOF 3.03xii |
| | Uptake of Influenza vaccine in persons 65+ years of age | Annual | 2014-15 | 70.2 | 71.4 | 69.2 | 72.7 | similar | | PHOF 3.03xiv |
| 25 | Priority Objective 4: Reducing Alcohol Harm | | 2012.11 | C1.1 | coc | | 645 | | | DUOS 2.40 |
| | Alcohol related admissions (ASR per 100,000 pop) Number of practitioners skilled in identifying those at risk from alcohol harm and delivering brief | Annual | 2013-14 April 2014 - | 614 | 606 | 541 | 645 | similar | - | PHOF 2.18 |
| | interventions (Local source) | Annual Local* | March 2015 | - | 500 | - | - | - | - | Lewisham Drugs & Alcohol Team |
| | Priority Objective 5: Preventing the uptake of smoking among children and young | | | umbers of p | | g | | | | |
| | Under 75 Mortality from Respiratory (DSR per 100,000 pop) | Annual | 2011-13 | 38.6 | 38.6 | 31.9 | 33.2 | similar | | PHOF 4.07i |
| | Under 75 Mortality from Lung Cancer (DSR per 100,000 pop) Smoking Prevalence(%) | Annual Annual | 2013 2013 | 31.3 21.4 | 46.9 20.6 | 31.0 17.3 | 33.7 18.4 | sig high similar | | NHS Indicator Portal - P00512 PHOf 2.14 |
| | 4 week smoking quitters (crude rate per 100,000) | Annual | 2014-15 | 751 | 680 | 531 | 522 | - | - | Smoking Quitters |
| | Number of 11-15 year-olds who take up smoking (%) | Every 2-3 years | 2010 | - | 9% | - | - | - | - | SHEU Survey (to be completed) |
| | Number of children in smoke free homes (%) | Every 2-3 years | 2010 | - | 57% | - | - | - | - | SHEU Survey (to be completed) |
| | Prevalence of Smoking in 15 year olds (proxy: % Never smoked at all - Yr8 and Yr10 children) Smoking status at time of delivery (%) | Every 2-3 years Annual/Qtr | 2010 2015-16 Q1 | 5.0 | 74% 5.0 | 5.0 | 10.7 | - | - | SHEU Survey (to be completed) HSCIC |
| | Priority Objective 6: Improving mental health and wellbeing | Aimadi) Qu | 2013 10 Q1 | 5.0 | | 5.0 | 20.7 | | | risere |
| 35 | Under 75 mortality rates for those with serious mental illness (DSR per 100,000 pop) | Annual | 2012-13 | 839.8 | 691.7 | - | 1,318.9 | sig low | - | NHSOF 1.5 |
| | Prevalence of Serious Mental Illness (%) | Annual | 2013-14 | 1.2 | 1.2 | 1.0 | 0.8 | - | | QOF |
| | Prevalence of Dementia (%) Prevalence of Depression (%) | Annual Annual | 2013-14 2013-14 | 0.3 5.3 | 0.4 4.8 | 0.4 4.4 | 0.6 6.5 | - | | QOF |
| - | Suicide rates (DSR per 100,000 pop) | Annual | 2013-14 | 7.5 | 8.1 | 7.2 | 8.8 | similar | * | PHOF 4.10 |
| 38 | Self-reported well-being - people with a low happiness score | Annual | 2013-14 | 10.2 | 8.1 | 9.6 | 9.7 | similar | | PHOF 2.23iii |
| | Priority Objective 7: Improving sexual health | | | 2.00 | 2=2: | | 407- | | | DUOS 2 CONTO CONTO CONTO |
| | Rate of chlamydia diagnoses per 100,000 young people aged 15 to 24 (crude rate) | Annual | 2014 | 3480 | 3504 | 2178 | 1978 | sig high | * | PHOF 3.02i/3.02ii (NCSP & CTAD) |
| | People presenting with HIV at a late stage of infection (%) Prevalence of diagnosed HIV infection per 1,000 among persons aged 15 to 59 years (crude rate) | Annual Annual | 2011-13 | 50.9 7.94 | 46.1 8.18 | 40.5 5.7 | 45.0 2.1 | similar | X | PHOF 3.04 PHE Sexual Health Profile |
| | Legal Abortion rate for all ages (crude rate per 1000 women) | Annual | 2013 | 26.6 | 25.0 | 21.8 | 16.5 | sig high | | ONS Abortion Stats |
| 42 | Teenage conceptions (Rate per 1,000 15-17 Yr olds) | Annual | 2013 | 42.0 | 33.1 | 21.8 | 24.3 | sig high | , J | PHE Sexual Health Profile |
| | Priority Objective 8 – Delaying and reducing the need for long term care and supp | | | | | | | | _ | |
| | Social care related quality of life (%) Rate of new admissions by older adults to long term care (crude rate per 100,000) | Annual Annual | 2013-14 2013-14 | 18.3 612.9 | 18.6 527.0 | 18.4 509.4 | 19 668.4 | - | | ASCOF 1A (P01507) ASCOF2A (P01514) (BCF) |
| 44 | % older people (65+) still at home 91 days after discharge from hospital into | | | | | | | | | |
| | rehabilitation/reablement services | Annual | 2013-14 | 86.5 | 86.9 | 88.9 | 81.9 | - | | ASCOF 2B |
| | Priority Objective 9: Reducing the number of emergency admissions for people w | th long term c | onditions | | | | | | | |
| 46 | Rate of avoidable emergency admissions - Unplanned Hospitalisation for chronic ambulatory care sensitive conditions (per 100,000 pop) | Annual/Qtr | 2014-15 | 953.2 | 1005.9 | 717.5 | 808.5 | sig high | • | CCGOI 2.6 |
| | % people able to manage effectively their own long term condition at home | Annual | 2013-14 | 61.1 | 58.5 | 59.7 | 64.4 | sig low | • | NHSOF 2.1 (BCF) |
| | Reviews of Adult Social Care Clients (cumulative % since Apr) | Annual/Qtr | 2014-15 Q3 | 0.7 | 63.7 | 0.7 | 0.7 | - | | BCF/Local Data - AO/D40 PPLUS |
| | Health-related quality of life for people with long-term conditions Emergency admissions for acute conditions that should not usually require hospital admission (DSR) | Annual | 2014-15 | 0.7 | 0.7 | 0.7 | 0.7 | - | | CCGOF 2.1 |
| 50 | rate per 100,000 Pop) | Annual | 2014-15 | 1294.5 | 1522.2 | 1047.8 | 1272.4 | sig high | 1 | CCGOF3.1 |
| 51 | Emergency Readmissions within 30 days of discharge (%) | Annual | 2014-15 | 15.69 | 15.54 | - | - | - | _ | Dr Foster |

Key

sig high -signficantly higher than England; sig low - significantly lower than England similar - statistically similar to England DSR - Directly Standardised Rates ASR - Age Standardised Rates ISR - Indirectly standardised Rates Lew - Lewisham; Lon - London; Eng - England

Links to Source with their abbreviations

http://www.phoutcomes.info/ http://www.phoutcomes.info/profile/sexualhealth https://www.indicators.ic.nhs.uk/webview/ http://www.hscic.gov.uk/qof http://ascof.hscic.gov.uk/

http://www.productivity.nhs.uk/ https://www.nhscomparators.nhs.uk/NHSComparators/HomePage.aspx

Note:

Annual/Qtr* - National Data available both quarterly and annually Annual* - Indicators not updated due to NO HES updates Qtr - Financial Quarters

*Indicator 51 refers to Lewisham CCG

Statistically Better than England
Statistically Similar to England
Statistically Worse than England
Blank where no statistical comparison could be made
Where performance is notably lower but other areas data is not s

Where performance is notably lower but other areas data is not statistically robust to compare

Arrows Indicate up or down performance of current year /qtr from previous yr/qtr


Public Health Outcomes Framework (PHOF)
Public Health England Sexual Health Profiles
NHS Indicator Portal (NHSIC) by Health and Social Care Information Centre (HSCIC)
Quality and Outcomes Framework(QOF) by HSCIC
Adult and Social Care Outcomes Framework (ASCOF)
NHS Better Care Better Value Indicators
NHS Comparators by HSCIC


Boroughs (Bromley, Bexley, Lambeth, Southwark, Greenwich and Lewisham) from London Bowel Screening Hub
Annual/Qtr** - Only Local Data available both quarterly and annually
Annual /Qtr*** - 2013/14 Q3 emergency admission rates are available on BCBV metrics for each Ambulator Care Sensitive (ACS) condition.
Local Ad-hoc - Bowel Screening data only available for all 6 South East London


Overarching Indicator


Health Outcome Indicator Rates


| | Target | 2006-08 | 2007-09 | 2008-10 | 2009-11 | 2010-12 | 2011-2013 | | | Direction of Travel from Target |
|--|--------|---------|---------|---------|---------|---------|-----------|-------------|-------------|---------------------------------------|
| Life Expectancy at Birth (Male)(yrs) | | 76.3 | 76.4 | 76.9 | 77.6 | 78.2 | 78.7 | 79.7 | 79.2 | |
| Life Expectancy at Birth (Female)(yrs) | | 81.2 | 81.6 | 81.7 | 82.3 | 82.6 | 83.0 | 83.8 | 83 | |
| Under 75 from CVD mortality (DSR) | | 132.3 | 119 | 110.3 | 96.7 | 91 | | 83.1 | 81.1 | |
| Slope index of inequality in life expectancy at birth within English local authorities, based on local deprivation deciles within each area (Male) | N/A | | | | 6 | 6.6 | | | | |
| Slope index of inequality in life expectancy at birth within English local authorities, based on local deprivation deciles within each area (Female) | N/A | | | | 6.3 | 6.6 | | | | |
| Infant Mortality rates | | | | | 4.8 | 4.9 | | 4.2 | 4.3 | |
| | | | | 2009-11 | 2010-12 | 2011-13 | | Lon 2011-13 | Eng 2011-13 | |
| Potential Years of Life Lost (PYLL) from causes considered amenable to healthcare (DSR) | | | | 2134.7 | 2102.0 | 2027.0 | | 1890.2 | 2023.5 | |
| | | 2008 | 2009 | 2010 | 2011 | 2012 | | Lon 2012 | Eng 2012 | |
| Low Birth Weight of all babies | | | | | 8.3 | 8.4 | | 7.9 | 7.3 | <u></u> |
| | | | | 2011/12 | 2012/13 | 2013/14 | | | | |
| Proportion of people using social care who receive self-directed support, and those receiving direct payments (Crude rate per 100,000) | | | | | 55.5 | 69.4 | | 67.5 | 62.1 | |
| Delayed transfers of care from hospital (crude rate per 100,000) | | | | 3.0 | 4.9 | | | 6.9 | 9.5 | 1 |
| Days of Delay (crude rate per 100,000) | | | | | 134 | | | | | |
| | | | | 2010 | 2011 | | | Lon 2011 | Eng 2011 | |
| Children in poverty (%) | | | | 31.7 | 30.5 | | | 26.5 | 20.6 | |

















Priority 1: Achieving Healthy Weight

| | Target | 2009/10 | 2010/11 | 2011/12 | 2012/13 | 2013/14 | London 2012/13 | England 2012/13 | Direction of Travel from Target |
|--|--------|---------|---------|---------|---------|---------|-------------------|--------------------|---------------------------------------|
| Excess weight in Adults (%) | | | | | 61.2 | | 57.3 | 63.8 | |
| Excess weight in Children - Reception Year (%) | | 27.8 | 24.8 | 24.8 | 25.0 | | 23.0 | 22.2 | |
| Excess Weight in Children- Year 6 (%) | | 40.6 | 39.1 | 40.4 | 38.3 | | 37.4 | 33.3 | |
| Breastfeeding Prevalence 6-8 weeks(%) | | | 74.6 | 75.7 | 73.5 | | 68.5 | 47.2 | |
| | | | 2010 | 2011 | 2012 | | London 2012 | England 2012 | |
| % of physically active and inactive adults - Active adults | | | | | 54.3 | | 57.2 | 56.0 | |
| % of physically active and inactive adults - Inactive adults | | | | | 29.2 | | 27.5 | 28.5 | |











Priority 2: Increasing the number of people who survive colorectal, breast and lung cancer for 1 and 5 years


| | Target | 2009 | 2010 | 2011 | 2012 | 2013 | London 2013 | | Direction of Travel from Target |
|--|--------|------------|------------|------------|------------|------------|--------------------------|---------------------|---------------------------------------|
| Cancer screening coverage - breast cancer (%) | | | 64.5 | 65.5 | 65.1 | 66 | 68.6 | 76.3 | |
| Cancer screening coverage - cervical cancer(%) | | | 71.7 | 71.4 | 71.9 | 72.4 | 68.6 | 73.9 | |
| Two week wait referrals (number per 100,000 population) | | | 1601 | 1810 | 1967 | 2273 | | 2166 | |
| | | 2012/13 Q4 | 2013/14 Q1 | 2013/14 Q2 | 2013/14 Q3 | 2013/14 Q4 | S.E London 2013/14 Q3 | • | |
| Cancer screening coverage - bowel cancer (%) | 60.0 | 44.2 | 44.2 | 38.2 | 35 | | 41.4 | 58.8 | |
| | | | 2010 | 2011 | 2012 | | London 2012 | England 2012 | |
| Early diagnosis of cancer (%) | | | | | 2273 | | | 2166 | |
| | | 2006-08 | 2007-09 | 2008-10 | 2009-11 | 2010-12 | London 2010-12 | England 2010- 12 | |
| Under 75 mortality from all cancers (DSR) | | 172.8 | 173.5 | 168 | 169.4 | 159.9 | 139.1 | 146.5 | |













Priority 3: Improving Immunisation Uptake


Percentage Uptake of Key Vaccines

| | Target | 2013/14 Q1 | 2013/14 Q2 | 2013/14 Q3 | 2013/14 Q4 | 2014/15 Q1 | London 2014/15 Q1 | England 2014/15 Q1 | Direction of Travel from Target |
|---|--------|------------|------------|------------|------------|------------|-------------------|--------------------|---------------------------------|
| | | | | | | | | | |
| Uptake of the first dose of Measles Mumps and Rubella vaccine (MMR1) at two | | | | | | | | | |
| years of age | 90.8% | 85.7% | 88.5% | 87.3% | 88.3% | 85.5% | 86.8% | 92.4% | 1 |
| Uptake of the third dose of Diptheria vaccine (D3) at one year of age | 91.9% | 86.7% | 88.4% | 88.3% | 88.9% | 90.0% | 88.6% | 93.9% | ₩ |
| Uptake of the fourth dose of Diphtheria vaccine (D4) at five years of age | 91.1% | 71.3% | 70.5% | 74.4% | 76.2% | 76.2% | 77.3% | 88.6% | |
| Uptake of the second dose of Measles Mumps and Rubella Vaccine (MMR2) at five | | | | | | | | | |
| years of age | 91.1% | 68.8% | 68.1% | 69.8% | 69.5% | 70.8% | 79.9% | 88.5% | . ↓ |
| | | | 2010/11 | 2011/12 | 2012/13 | 2013/14 | London 2012/13 | England 2012/13 | |
| | | | | | | | | | |
| Uptake of Human Papilloma Virus (HPV) vaccine in girls in Year 8 in Lewisham | | | | | | | | | |
| Schools | | | 68.8 | 80.6 | 84.8 | | 78.9 | 86.1 | ☆ |
| Uptake of Influenza vaccine in those over 65 years of age | | | 69.6% | 70.1% | 68.2% | | 71.2% | 73.4% | 企 |

Source: HPA Cover Data








Priority 4: Reducing Alcohol Harm


| | Target | 2009/10 | 2010/11 | 2011/12 | 2012/13 | 2013/14 | London 2012/13 | | Direction of Travel from Target |
|--|--------|---------|---------|---------|---------|---------|-------------------|-----|---------------------------------------|
| Alcohol related admissions (DSR) | - | - | - | 588 | 614 | - | 554 | 637 | |
| Number of practitioners skilled in identifying those at risk from alcohol harm and delivering brief interventions (Local source) | - | - | - | - | 120 | - | | | |
| | | | | | | | | | |





Priority 5: Preventing the uptake of smoking among children and young people and reducing the numbers of people smoking

Tobacco Control

| | Target | 2006-08 | 2007-09 | 2008-10 | 2009-11 | 2010-12 | | England 2010- 12 | Direction of Travel from Target |
|--|--------|------------|------------|------------|------------|---------|----------------|---------------------|---------------------------------------|
| Under 75 Mortality from Respiratory | | 51.3 | 49.7 | 47 | 40.9 | 38.6 | 32.6 | 33.5 | |
| | | 2008 | 2009 | 2010 | 2011 | 2012 | London 2012 | England 2012 | |
| Under 75 Mortality from Lung Cancer | | 30.39 | 24.92 | 26.68 | 35.78 | 21.0 | 23.04 | 24.06 | |
| Smoking Prevalence (18+) - routine and manual | | | | | 25.4 | 24.3 | 25.7 | 29.7 | |
| | | 2013/14 Q1 | 2013/14 Q2 | 2013/14 Q3 | 2013/14 Q4 | | | England 2013/14 | |
| 4 week smoking quitter (crude rate per 100,000) | | 200.3 | 197.4 | 197.3 | 180.7 | 172.6 | 656 | 688 | |
| Smoking at time of delivery | | 5.9% | 4.9% | 5.3% | 5.2% | 5.4% | | 11.8% | |
| | | | | 2008 | 2010 | | | | |
| Number of 11-15 year-olds who take up smoking (%) | | | | 16.0% | 9.0% | | | | |
| Number of children in smoke free homes (%) | | | | 56.0% | 57.0% | | | | |
| Prevalence of Smoking in 15 year olds (proxy: % Never smoked at all - Yr8 and Yr10 children) | | | | 95.0% | 74.0% | | | | |











Priority 6: Improving mental health and wellbeing


| | Target | 2009/10 | 2010/11 | 2011/12 | 2012/13 | 2013/14 | London 2011/12 | | Direction of Travel from Target |
|--|--------|---------|---------|---------|---------|---------|-------------------|---------------|---------------------------------------|
| Under 75 mortality rates for those with serious mental illness (DSR) | | 302.2 | 303.6 | 305.3 | | | - | 337.4 | |
| Prevalence of SMI | | | | 1.2 | 1.2 | | 1.0 | 0.8 | |
| Prevalence of Dementia | | 0.3 | 0.3 | 0.3 | 0.3 | | 0.4 | 0.6 | |
| Prevalence of Depression | | 10 | 10.1 | 10.4 | 5.3 | | 4.4 | 5.8 | |
| Self-reported well-being - people with a low happiness score | | | | 15 | 10.2 | | 10.3 | 10.4 | |
| | | 2007-09 | 2008-10 | 2009-11 | 2010-12 | | London | England 2010- | |
| | | | | | | | 2010-12 | 12 | |
| Suicide rates | | 6.9 | 7 | 7.1 | 7.5 | | 7.5 | 8.5 | |












Priority 7: Improving sexual health


| | Target | 2009 | 2010 | 2011 | 2012 | 2013 | London 2013 | England 2013 | Direction of Travel from |
|--|--------|------|------|---------|---------|------|----------------|---------------|-----------------------------|
| ** old data - 2012 | | | | | | | | | Target |
| Rate of chlamydia diagnoses per 100,000 young people aged 15 to 24 (crude rate) | | | 4728 | 4762 | 4179 | 3480 | 2179 | 2016 | |
| Prevalence of diagnosed HIV infection per 1,000 among persons aged 15 to 59 years (crude rate)** | | | 7.5 | 7.8 | 7.9 | | 5.5 | 2.1 | |
| Legal Abortion rate for all ages (crude rate per 1000 women) | | 32.3 | 29.9 | 32.3 | 27.4 | 26.6 | 21.7 | 16.1 | |
| Teenage conceptions** | | 48.5 | 41.8 | 39.9 | 42.0 | | 25.9 | 27.7 | |
| | | | | 2009-11 | 2010-12 | | London | England 2010- | |
| | | | | | | | 2010-12 | 12 | |
| People presenting with HIV at a late stage of infection(%) | | | | 51.8 | 50.9 | | 44.9 | 48.3 | |
| | | | | | | | | | |


Trend Charts


60.0 50.0 40.0 30.0


0.0 crude rate 0.0














Priority 8: Delaying and reducing the need for long term care and support.

| | Target | 2009/10 | 2010/11 | 2011/12 | 2012/13 | 2013/14 | London 2012/13 | | Direction of Travel from Target |
|--|--------|---------|---------|---------|---------|---------|-------------------|-------|---------------------------------------|
| social care related quality of life (%) | 19 | | 17.7 | 17.9 | 18.3 | 18.6 | 18.4 | 19 | |
| Rate of new 65+ admissions to long term care (crude rate per 100,0 | 549.4 | 931.2 | 854.3 | 560.7 | 612.9 | 527 | 509.4 | 668.4 | |
| % older people (65+) still at home 91 days after discharge from hospital into rehabilitation/reablement services | 88.0 | 88.2 | 88.6 | 89.4 | 86.5 | 86.9 | 88.9 | 81.9 | |








Priority 9: Reducing the number of emergency admissions for people with long term conditions


| | Target | 2009/10 | 2010/11 | 2011/12 | 2012/13 | 013/14 (Prov | London 2013/14 (Prov) | England 2013/14 (Prov) | Direction of Travel from Target |
|--|--------|---------|---------|------------------------|------------------------|------------------------|----------------------------------|-----------------------------------|---------------------------------------|
| Rate of avoidable emergency admissions (Std rate per 100,000 pop) | | 14.52 | 1061.2 | 1029 | 1027.6 | 940.9 | 734.6 | 780.9 | |
| | | | | | 2012/13 | 2013/14 | London 2013/14 | England 2013/14 | |
| % people able to manage effectively their own long term condition at home | | | | 58.1 | 62.3 | 61.1 | 59.7 | 65.1 | |
| | | | | | | | London 2012/13 | England 2012/13 | |
| Reviews of Adult Social Care Clients (%) | 70 | 74.7 | 62 | 64.4 | 77.9 | 70.7 | 69.8 | 66.6 | |
| | | | | | | | | England 2011/12 | |
| Emergency Readmissions within 30 days of discharge from hospital | | 12.11 | 11.96 | 12.73 | | | | 11.78 | |
| | | | | Jul 2011 - Mar 2012 | Jul 2012 - Mar 2013 | Jul 2013 - Mar 2014 | London Jul 2012 - Mar 2013 | England Jul 2012 - Mar 2013 | |
| Health-related quality of life for people with long-term conditions | | | | 0.73 | 0.75 | 0.75 | 0.75 | 0.74 | |
| | | | 2010/11 | 2011/12 | 2012/13 | 2013/14 (Prov) | London 2013/14 | England 2013/14 | |
| Emergency admissions for acute conditions that should not usually require hospital admission | | | 1147.8 | 1155.8 | 1324.8 | 1279.4 | 991 | 1164.7 | |

| | | 2010/11 | 2011/12 | 2012/13 | 2013/14(Pr ov) | London 2013/14 Prov | England 2013/14 Prov | |
|---|--|---------|---------|---------|-------------------|---------------------------|-------------------------|--|
| Unplanned admission of chronic ambulatory care sensitive conditions | | 1061.2 | 1029.0 | 1027.6 | 940.9 | 734.6 | 780.9 | |

Trend Charts



Emergency Readmissions within 30 days of discharge from hospital



100 80 60 40 20 0 2011/12 2012/13 London 2013/14 England 2013/14 (Prov) (Prov)

Unplanned admission of chronic ambulatory care

Emergency admissions for acute conditions that should

Reviews of Adult Social Care Clients (%)

